



100 North Wren Drive • Pittsburgh, PA 15243 • Phone (412) 429-2570 • Fax (412) 429-2572  
95 West Beau Street • Washington, PA 15301 • Phone (412) 429-2570 • Fax (724) 228-8822  
521 East Bruceton Road • Pittsburgh, PA 15236 • Phone (412) 429-2570 • Fax (412) 714-4591

**PRESCRIPTION NAME:** Antifungal Cream (Ciclopirox 3%, itraconazole 5%, urea 20%)

**COMMON USES:** This cream is used to treat fungal infections of the skin

**BEFORE USING THIS MEDICATION:** Tell your provider if you have an allergy to ciclopirox, itraconazole, urea, or any other part of this drug. Tell your provider about the allergy and what symptoms you had (rash, hives, itching, shortness of breath, wheezing, cough, swelling of face, lips, tongue or throat or any other signs).

**WHAT DOES THIS MEDICATION DO?:** This medication contains 3 active ingredients. Ciclopirox and itraconazole are antifungal/antimicrobial medications to kill fungus. Urea helps decrease scaling and firmness of treated areas.

**HOW TO APPLY THIS MEDICATION:** Use this medication as instructed by your provider. Cleanse skin prior to application of cream and pat dry skin. Apply a thin layer of lotion on the affected areas of skin twice a day, or as directed by your provider.

**COMMON SIDE EFFECTS:** May experience minor burning, itching dryness, redness with use of this medication. Use of other skin products while using this medication may cause more irritation.

**IMPORTANT INFORMATION:**

- Advised your provider of all medication you are taking to avoid any reactions.
- If you have an allergy to ciclopirox, itraconazole, urea or other part of this medication, tell your provider.
- Stop application if you become pregnant or are breastfeeding. Use birth control while taking this medication and for at least 2 months after your last dose.
- Do not swallow this medication. If swallowed call provider or poison control center immediately.

**SERIOUS SIDE EFFECTS:** Even though it may be rare, some patients may have very bad and sometimes serious side effects when taking a medication. Tell your provider if you have any of the following signs or symptoms of an allergic reaction such as a rash, hives, swelling, blistering, wheezing, tightness in the chest or throat, trouble breathing or swallowing, unusual hoarseness, swelling of the mouth, face, lips, tongue, or throat. Rare serious side effects such as heart failure (weak heart), drug-drug interactions, high blood sugars, high blood pressure have occurred with oral itraconazole. Symptoms of heart failure include shortness of breath, weight gain, coughing, fast heartbeat, swelling in the arms/legs. Do not take this medication if you are taking any of these drugs: avanafil, cisapride, dihydroergotamine, disopyramide, dofetilide, dronedarone, eplerenone, ergonovine, ergotamine, felodipine, irinotecan, isavuconazole, ivabradine, lomitapide, lovastatin, lurasidone, methadone, methylergonovine, oral midazolam, naloxegol, nisoldipine, pimozide, quinidine, ranolazine, simvastatin, ticagrelor, trazolam. If you have kidney or liver problems, do not take this medication if you are also taking fesoterodine, solifenacin, or colchicine. Signs of electrolyte problems include mood changes, confusion, muscle pain, weakness, abnormal heartbeat, low hunger, vomiting. Signs of high blood sugars include confusion, sleepiness, more thirst, more hunger, passing urine more often, flushing, fast breathing, breath that smells like fruit. Signs of high or low blood pressure includes very bad headache, passing out, change in eyesight, chest pain. Other rare side effects of oral itraconazole includes ringing in the ears, hearing loss, swelling, lowered interest in sex, difficulty in getting or keeping an erection, fever or chills, liver problems, headache, vomiting, fatigue.

If any skin reactions happen, let your provider know. If you have any questions, please call our office at 412-429-2570.